


PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : C07D 241/04, 243/08, 295/192, 403/06, A61K 31/495, 31/55</p>	<p>A1</p>	<p>(11) International Publication Number: WO 98/20001 (43) International Publication Date: 14 May 1998 (14.05.98)</p>
<p>(21) International Application Number: PCT/US97/20020 (22) International Filing Date: 4 November 1997 (04.11.97) (30) Priority Data: 60/029,894 8 November 1996 (08.11.96) US (71) Applicant: BRISTOL-MYERS SQUIBB COMPANY [US/US]; P.O. Box 4000, Princeton, NJ 08543-4000 (US). (72) Inventor: LEFTHERIS, Katherina; 92 Richmond Drive, Skillman, NJ 08558 (US). (74) Agents: HOFFMAN, Frank, P. et al.; Bristol-Myers Squibb Company, P.O. Box 4000, Princeton, NJ 08543-4000 (US).</p>		<p>(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report.</i></p>
<p>(54) Title: THIOPROLINE-CONTAINING INHIBITORS OF FARNESYL PROTEIN TRANSFERASE</p> <p>(57) Abstract</p> <p>Inhibition of farnesyl transferase, which is an enzyme involved in ras oncogene expression, is effected by compounds of formula (I) its enantiomers, diastereomers, and pharmaceutically acceptable salts, prodrugs and solvates.</p> <div style="text-align: center;">  <p>(I)</p> </div>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

THIOPROLINE-CONTAINING INHIBITORS OF FARNESYL PROTEIN TRANSFERASE

Field of the Invention

5 This invention relates to compounds that inhibit farnesyl-
protein transferase and Ras protein farnesylation, thereby making them
useful as anti-cancer agents. The compounds are also useful in the
treatment of diseases, other than cancer, associated with signal
transduction pathways operating through Ras and those associated with
10 CAAX-containing proteins other than Ras that are also post-
translationally modified by the enzyme farnesyl protein transferase.
The compounds may also act as inhibitors of other prenyl transferases,
and thus be effective in the treatment of diseases associated with other
prenyl modifications of proteins.

15

Background of the Invention


The mammalian ras gene family comprises three genes: H-ras, K-
ras and N-ras. The Ras proteins are a family of GTP-binding and
hydrolyzing proteins that regulate cell growth and differentiation.
20 Overproduction of normal Ras proteins or mutations that inhibit their
GTPase activity can lead to uncontrolled cell division.

The transforming activity of Ras is dependent upon localization of
the protein to plasma membranes. This membrane binding occurs via a
series of post-translational modifications of the cytosolic Ras proteins. The
25 first and mandatory step in this sequence of events is the farnesylation of
these proteins. The reaction is catalyzed by the enzyme farnesyl protein
transferase (FPT), and farnesyl pyrophosphate (FPP) serves as the farnesyl
group donor in this reaction. The Ras C-terminus contains a sequence
motif termed a "Cys-Aaa1-Aaa2-Xaa" box (CAAX box), wherein Cys is
30 cysteine, Aaa is an aliphatic amino acid, and Xaa is a serine or
methionine. Farnesylation occurs on the cysteinyl residue of the CAAX
box (Cys-186), thereby attaching the prenyl group on the protein via a thio-
ether linkage.

Brief Description of the Invention


In accordance with the present invention, a compound of the formula

5 I



its enantiomers and diastereomers, and pharmaceutically acceptable salts, prodrugs and solvates thereof inhibit S-farnesyl protein transferase which is an enzyme involved in Ras oncogene function. In formula I and throughout this specification, unless otherwise specified, the symbols are defined as follows:

A is



optionally substituted, at any available position or positions on the ring, with one or more of the following substituents, halo, alkyl, alkoxy, aryl, hydroxy, alkanoyl, alkanoyloxy, amino, alkylamino, dialkylamino, alkanoylamino, thiol, alkylthio, alkylthiono, alkylsulfonyl, sulfonamido, nitro, cyano, carboxy, carbamyl or alkoxycarbonyl;

B is $-\text{CH}_2-$ or $-\text{C}(\text{O})-$;

20 B² is alkyl, aryl or heterocycle;

V and W are each, independently, $-\text{CH}_2-$ or $-\text{C}(\text{O})-$;

X is $-\text{SH}$, $-\text{OH}$ or $-\text{NHR}^6$;

X' is $-\text{NR}^7-$, $-\text{CH}_2-$ or $-\text{CH}(\text{NHR}^8)-$;

R, R¹, R² and R³ are each independently, hydrogen, alkyl, alkenyl, alkynyl, aryl, heterocycle, $-\text{C}(\text{O})\text{NR}^4\text{R}^5$, or $-\text{C}(\text{O})\text{OR}^4$; or any two of R, R¹, R² and R³, if present, may also be alkylene attached to the same carbon atom forming a spiro ring;

R^4 and R^5 are each independently selected from hydrogen, hydroxyl, alkyl, cycloalkyl, heterocycle, aryl, aroyl, heteroaroyl, arylsulfonyl or heteroarylsulfonyl; or R^4 and R^5 may together form a 5- to 7-membered saturated ring with the atom to which they are attached;
5 said ring being optionally substituted with halogen, hydroxy, alkoxy, oxo, nitro, cyano, $-C(O)H$, $-C(O)OH$ and the like;

R^6 , R^7 and R^8 are each independently, hydrogen or alkyl;

m is an integer from 0 to 1; and

n is an integer from 1 to 2.

10

Detailed Description of the Invention

Definition of Terms

Listed below are definitions of various terms used to describe this invention. These definitions apply to the terms as they are used throughout this specification, unless otherwise limited in specific instances, either individually or as part of a larger group.

The term "alkyl" refers to optionally substituted straight or branched chain hydrocarbon groups having 1 to 20 carbon atoms, preferably 1 to 8 carbon atoms. Exemplary unsubstituted alkyl groups include methyl, ethyl, propyl, isopropyl, n-butyl, t-butyl, isobutyl, pentyl, hexyl, isohexyl, heptyl, 4,4-dimethylpentyl, octyl and the like. The term "alkyl" also includes substituted alkyl groups. Substituted alkyl groups refer to alkyl groups substituted by one or more of the following groups: halo (such as CCl_3 or CF_3), hydroxy, -OR¹¹, cycloalkyl, alkanoyl, alkanoyloxy, amino, alkylamino, dialkylamino, alkanoylamino, thiol, alkylthio, alkylthiono, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, sulfonamido, nitro, cyano, carboxy, carbamyl, alkoxycarbonyl, aryl, aralkoxy, phenyl, substituted phenyl, guanidino, indolyl, imidazolyl, furyl, thienyl, thiazolyl, pyrrolidyl, pyridyl, pyrimidyl and the like.

The term "lower alkyl" refers to those alkyl groups as described above having 1 to 4 carbon atoms.

The term "halogen" or "halo" refers to fluorine, chlorine, bromine and iodine.

The term "alkenyl" refers to any of the above alkyl groups having at least 2 carbon atoms and further containing at least one carbon to carbon double bond. Groups having two to four carbon atoms are preferred.

The term "alkynyl" refers to any of the above alkyl groups having at least two carbon atoms and further containing at least one carbon to carbon triple bond. Groups having two to four carbon atoms are preferred.

The term "alkylene" refers to a straight chain bridge of 1 to 6 carbon atoms connected by single bonds (e.g., $-(CH_2)_x-$ wherein x is 1 to 6), which may be substituted with 1 to 3 lower alkyl groups.

5 The term "cycloalkyl" refers to cyclic hydrocarbon groups of 3 to 8 carbon atoms.

The term "alkoxy" refers to alkyl-O-.

The term "alkanoyl" refers to alkyl-C(O)-.

The term "alkanoyloxy" refers to alkyl-C(O)-O-.

10 The terms "alkylamino" and "dialkylamino" refer to (alkyl)NH- and (alkyl)₂N-, respectively.

The term "alkanoylamino" refers to alkyl-C(O)-NH-.

The term "alkylthio" refers to alkyl-S-.

The term "alkylthiono" refers to alkyl-S(O)-.

The term "alkylsulfonyl" refers to alkyl-S(O)₂-.

15 The term "carbamyl" refers to -C(O)NR⁴R⁵.

The term "alkoxycarbonyl" refers to alkyl-O-C(O)-.

The term "aryl" refers to monocyclic or bicyclic aromatic hydrocarbon groups having 6 to 12 carbon atoms in the ring portion, such as phenyl, naphthyl, tetrahydronaphthyl, biphenyl and diphenyl
20 groups, each of which may optionally be substituted by one to four substituents such as alkyl, halo, hydroxy, alkoxy, alkanoyl, alkanoyloxy, amino, alkylamino, dialkylamino, alkanoylamino, thiol, alkylthio, nitro, cyano, carboxy, carboxyalkyl, carbamyl, alkoxycarbonyl, alkylthiono, alkylsulfonyl, sulfonamido, heterocyclo and the like.

25 The term "aralkyl" refers to an aryl group bonded directly through an alkyl group, such as benzyl.

The term "aralkoxy" refers to an aryl group bonded directly through an alkoxy group.

The term "arylsulfonyl" refers to aryl-S(O)₂-.

30 The term "aroyl" refers to aryl-C(O)-.

The term "heterocycle" refers to an optionally substituted, fully saturated or unsaturated, aromatic or nonaromatic cyclic group, for example, which is a 4 to 7 membered monocyclic, 7 to 11 membered

bicyclic, or 10 to 15 membered tricyclic ring system, which has at least one heteroatom in at least one carbon atom-containing ring. Each ring of the heterocyclic group containing a heteroatom may have 1, 2 or 3 heteroatoms selected from nitrogen atoms, oxygen atoms and sulfur atoms, where the nitrogen and sulfur heteroatoms may also optionally be oxidized and the nitrogen heteroatoms may also optionally be quaternized. The heterocyclic group may be attached at any heteroatom or carbon atom.

Exemplary monocyclic heterocyclic groups include pyrrolidinyl, pyrrolyl, pyrazolyl, oxetanyl, pyrazolinyl, imidazolyl, imidazolinyl, imidazolidinyl, oxazolyl, oxazolidinyl, isoxazolinyl, isoxazolyl, thiazolyl, thiadiazolyl, thiazolidinyl, isothiazolyl, isothiazolidinyl, furyl, tetrahydrofuryl, thienyl, oxadiazolyl, piperidinyl, piperazinyl, 2-oxopiperazinyl, 2-oxopiperidinyl, 2-oxopyrrolodiny, 2-oxoazepinyl, azepinyl, 4-piperidonyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, tetrahydropyranyl, morpholinyl, thiamorpholinyl, thiamorpholinyl sulfoxide, thiamorpholinyl sulfone, 1,3-dioxolane and tetrahydro-1,1-dioxothienyl, and the like.

Exemplary bicyclic heterocyclic groups include indolyl, benzothiazolyl, benzoxazolyl, benzothienyl, quinuclidinyl, quinolinyl, tetra-hydroisoquinolinyl, isoquinolinyl, benzimidazolyl, benzopyranyl, indolizinyl, benzofuryl, chromonyl, coumarinyl, benzopyranyl, cinnolinyl, quinoxalinyl, indazolyl, pyrrolopyridyl, furopyridinyl (such as furo[2,3-c]pyridinyl, furo[3,2-b]pyridinyl] or furo[2,3-b]pyridinyl), dihydroisoindolyl, dihydroquinazolinyl (such as 3,4-dihydro-4-oxo-quinazolinyl) and the like.

Exemplary tricyclic heterocyclic groups include carbazolyl, benzidolyl, phenanthrolinyl, acridinyl, phenanthridinyl, xanthenyl and the like.

The term "heterocycle" also includes substituted heterocycle groups. Substituted heterocycle groups refer to heterocycle groups substituted with 1, 2 or 3 of the following:

- (a) alkyl;
- 5 (b) hydroxy (or protected hydroxy);
- (c) halo;
- (d) oxo (i.e. = O);
- (e) amino, alkylamino or dialkylamino;
- (f) alkoxy;
- 10 (g) cycloalkyl;
- (h) carboxy;
- (i) heterocyclooxy;
- (j) alkoxycarbonyl, such as unsubstituted lower
alkoxycarbonyl;
- 15 (k). carbamyl, alkylcarbamyl dialkylcarbamyl;
- (l) mercapto;
- (m) nitro;
- (n) cyano;
- (o) sulfonamido, sulfonamidoalkyl or
20 sulfonamidodialkyl;
- (p) aryl;
- (q) alkylcarbonyloxy;
- (r) arylcarbonyloxy;
- (s) arylthio;
- 25 (t) aryloxy;
- (u) alkylthio;
- (v) formyl;
- (w) arylalkyl; or
- (x) aryl substituted with alkyl, cycloalkyl,
30 alkoxy, hydroxy, amino, alkylamino,
dialkylamino or halo.

The term "heterocyclooxy" denotes a heterocyclic group bonded through an oxygen bridge.

The term "heteroaryl" refers to an aromatic heterocycle.

The term "heteroarylsulfonyl" refers to heteroaryl-S(O)₂-.

5 The term "heteroaroyl" refers to heterocyclo-C(O)-.

Throughout the specification, groups and substituents thereof are chosen to provide stable moieties and compounds.

The compounds of formula I form salts which are also within the scope of this invention. Pharmaceutically acceptable (i.e., non-toxic,
10 physiologically acceptable) salts are preferred, although other salts are also useful, e.g., in isolating or purifying the compounds of this invention.

The compounds of formula I may form salts with alkali metals such as sodium, potassium and lithium, with alkaline earth metals
15 such as calcium and magnesium, with organic bases such as dicyclohexylamine, tributylamine, pyridine and amino acids such as arginine, lysine and the like. Such salts may be obtained, for example, by exchanging the carboxylic acid protons, if they contain a carboxylic acid, in compound I with the desired ion in a medium in which the salt
20 precipitates or in an aqueous medium followed by evaporation. Other salts can be formed as known to those having ordinary skill in the art.

The compounds of formula I may form salts with a variety of organic and inorganic acids. Such salts include those formed with hydrogen chloride, hydrogen bromide, methanesulfonic acid, sulfuric
25 acid, acetic acid, trifluoroacetic acid, maleic acid, benzenesulfonic acid, toluenesulfonic acid and various others (e.g., nitrates, phosphates, borates, tartrates, citrates, succinates, benzoates, ascorbates, salicylates and the like). Such salts may be formed by reacting compound I in an equivalent amount of the acid in a medium in which the salt precipitates
30 or in an aqueous medium followed by evaporation.

In addition, zwitterions ("inner salts") may be formed.

A compound of the formula I may also have prodrug forms. Any compound that will be converted *in vivo* to provide the bioactive agent (i.e., the compound of formula I) is a prodrug within the scope and spirit of the invention.

5 Various forms of prodrugs are well known in the art. For examples of such prodrug derivatives, see:

- a) Anne H. Kahns and Hans Bundgaard, *International Journal of Pharmaceutics*, 62 (1990), p. 193-205.
- b) *Design of Prodrugs*, edited by H. Bundgaard, (Elsevier, 1985) and *Methods in Enzymology*, Vol. 42, 309-396, edited by K. Widder et al. (Academic Press, 1985);
- c) *A Textbook of Drug Design and Development*, edited by Krogsgaard-Larsen and H. Bundgaard, Chapter 5, "Design and Application of Prodrugs," by H. Bundgaard, 113-191
10 (1991);
- d) H. Bundgaard, *Advanced Drug Delivery Reviews*, 8, 1-38 (1992);
- e) H. Bundgaard et al., *Journal of Pharmaceutical Sciences*, 77, 285 (1988); and
- f) N. Kakeya et al., *Chem Pharm Bull*, 32, 692 (1984).
15 20

It should further be understood that solvates (e.g., hydrates) of the compounds of formula I are also within the scope of the present invention. Methods of solvation are generally known in the art.

25 Use and Utility

The compounds of formula I are inhibitors of S-farnesyl protein transferase. They are thus useful in the treatment of a variety of cancers, including (but not limited to) the following:

- carcinoma, including that of the bladder, breast,
30 colon, rectum, kidney, liver, lung, ovary, pancreas,
 stomach, cervix, thyroid and skin;
 hematopoietic tumors of lymphoid lineage, including
 acute lymphocytic leukemia, B-cell lymphoma and Burkitts
 lymphoma;

hematopoietic tumors of myeloid lineage, including
acute and chronic myelogenous leukemias and
promyelocytic leukemia;

5 tumors of mesenchymal origin, including
fibrosarcoma and rhabdomyosarcoma;
tumors of the central and peripheral nervous system;
and

other tumors, including melanoma, seminoma,
tetratocarcinoma, neuroblastoma, astrocytoma and glioma.

10 The compounds of formula I are especially useful in treatment of
tumors having a high incidence of ras involvement, such as colon, lung,
and pancreatic tumors; as well as tumors adversely affected by
inhibition of farnesyl transferase or prenyl transferase. By the
administration of a composition having one (or a combination) of the
15 compounds of this invention, development of or progression of tumors in
a mammalian host is reduced.

Compounds of formula I may also be useful in the treatment of
diseases other than cancer that may be associated with signal
transduction pathways operating through ras, e.g., neuro-fibromatosis,
20 atherosclerosis, pulmonary fibrosis, arthritis, psoriasis,
glomerulonephritis, polycystic kidney disease, restenosis following
angioplasty or vascular surgery, hypertrophic scar formation, endotoxic
shock and autoimmune and inflammatory diseases such as rheumatoid
arthritis, asthma, urticaria, angioedema, systemic sclerosis and
25 scleroderma, vasculitis, gout, inflammatory lung disease (emphysema,
bronchitis, fibrosis), adult respiratory distress syndrome (ARDS), Lupus
erythematosus, arteriosclerosis, inflammatory bowel disease (Crohn's
disease, ulcerative colitis), periodontal disease, multiple sclerosis,
myasthenia gravis, graft vs. host disease, transplantation rejection,
30 Sjogren's disease, Graves disease and type 1 diabetes. Compounds of
formula I may be useful as anti-fungal agents.

Compounds of formula I may also be useful in the treatment of diseases associated with farnesyl transferase substrates other than ras (e.g., nuclear lamins and transducin) that are also post-translationally modified by the enzyme farnesyl protein transferase.

5 Compounds of I may also act as inhibitors of other prenyl transferases (e.g., geranylgeranyl transferase I and II), and thus be effective in the treatment of diseases associated with other prenyl modifications (e.g., geranylgeranylation) of proteins (e.g., the rap, rab, rac and rho gene products and the like). For example, they may find use
10 as drugs against Hepatitis delta virus (HDV) infections, as suggested by the recent finding that geranylgeranylation of the large isoform of the delta antigen of HDV is a requirement for productive viral infection (J.S. Glen et al., *Science*, 256, 1331 (1992)).


The compounds of this invention may also be useful in
15 combination with known anti-cancer and cytotoxic agents and treatments, including radiation. If formulated as a fixed dose, such combination products employ the compounds of this invention within the dosage range described below and the other pharmaceutically active agent within its approved dosage range. Compounds of formula I may
20 be used sequentially with known anticancer or cytotoxic agents and treatments, including radiation when a combination formulation is inappropriate.

Farnesyl transferase assays were performed as described in V. Manne et al., *Drug Development Research*, 34, 121-137 (1995). The
25 compounds of Examples 1-8 inhibited farnesyl transferase with IC₅₀ values between 0.1nM and 10mM.

The compounds of this invention may be formulated with a pharmaceutical vehicle or diluent for oral, parenteral, intravenous, subcutaneous, rectal, vaginal or topical administration. The pharmaceutical composition can be formulated in a classical manner using solid or liquid vehicles, diluents and additives appropriate to the desired mode of administration. Orally, the compounds can be administered in the form of tablets, capsules, granules, powders and the like. The compounds may be administered in a dosage range of about 0.05 to 50 mg/kg/day, preferably less than 50 mg/kg/day, in a single dose or in 2 to 4 divided doses.


Compounds of formula I where B is CH₂ may be prepared by coupling compounds of formula

II




where Y is a protected thiol, protected alcohol or protected amine with compounds of formula

III



by reductive amination using a reducing agent such as sodium triacetoxyborohydride or sodium cyano borohydride to form compounds of formula

IV



Compounds of formula IV are deprotected to provide compounds of formula I where B is CH₂ using standard methods known in the literature (*Protecting Groups in Organic Synthesis*, 2nd Edition, Theodora Green, Peter Wuts, John Wiley & Sons, Inc. 1991).


- 5 Alternatively compounds of formula IV where Y is a trityl S- can be deprotected using acidic conditions such as 50% trifluoroacetic acid in methylene chloride in the presence of a scavenger such as triethylsilane.

Alternatively, compounds of formula I where X is OH or -NHR⁶ may also be prepared by reductive amination using compounds of
10 formula II and III and hydrogen with a catalyst such as palladium on carbon or palladium hydroxide on carbon.


Compounds of formula I where B is C(O) can be prepared by coupling compounds of formula

V

15



with a compound of formula III in the presence of a coupling agent (e.g., dicyclohexylcarbodiimide (DCC)) to form compounds of formula
VI




20

Compounds of formula VI are converted to compounds of formula I where B is C(O) by deprotecting using standard methods found in the literature.


Compounds of formula II and V are prepared by methods known in
25 the art. See, for example, V. Eskwarakrishnan and L. Field, *J. Org. Chem.*, 46, 4182 (1981); D. Papaioannou et al., *Acta Chemica Scandinavia*, 44, 243 (1990); A. Meyers, *Org. Syn.*, 51, 103 (1971); T. Rosen et al.,

Synthesis, 40, (1988); C. Agami et al., *Tetrahedron*, 48, 431 (1992); Y. Ueda and V. Vinet, *Can. J. Chem.*, 64, 2184 (1986); Sinagawa et al., *J. Antibiotico*, 43 (1990) p. 519-532; and EP 696593-A2.

- Compounds of formula III where V and W are $-\text{CH}_2-$, $m=0$, B^2 is aryl, R^1 , R^2 and R^3 are each hydrogen, R is attached to V and is alkyl (other than arylsulfonyl or heteroarylsulfonyl substituted alkyl) are prepared by coupling a compound of formula VII




- 10 wherein R^9 is alkyl, arylalkyl or aryl and Prot^1 is an amine protecting group (e.g., t-butyloxycarbonyl (BOC), benzyloxycarbonyl (Cbz) and the like): in the presence of a coupling agent (e.g., dicyclohexylcarbodiimide (DCC)) with a carboxylic acid derivative of the formula VIII



15


wherein R^{10} is alkyl, arylalkyl or aryl, to form a compound of the formula

IX



5 Compounds of formula IX can then be treated with a suitable N-deprotecting agent (e.g., HCl for BOC) and cyclized to provide a compound of formula


X



10

A compound of formula X can be reduced with a reducing agent (e.g., lithium aluminum hydride (LiAlH_4)) to form the compound of the formula


XI



15

The compound of formula XI can then be protected with a protecting agent (e.g., di-*t*-butyloxy dicarbonate) to form a compound of the formula

XII



wherein Prot² is a suitable protecting group, (e.g. t-butyloxycarbonyl).


- 5 The hydroxy group in the compound of formula XII can then be alkylated under basic conditions (e.g., sodium hydride with a compound of formula

XIIa



- 10 where L is a leaving group, for example a halide or sulfonyl ester; and R¹¹ is alkyl, alkenyl, alkynyl, cycloalkyl, aryl, aroyl, heteroaroyl, arylsulfonyl, heteroarylsulfonyl, heterocycle or -C(O)NR⁴R⁵) to form a compound of the formula


XIII



15

Compounds of the formula XIII can then be hydrogenated using a catalyst in the presence of hydrogen (e.g., palladium/carbon/hydrogen) to form a compound of the formula


XIV




A compound of formula XIV can then be coupled with a compound
 5 of the formula
 XIVa



wherein Ar is an aryl group, in the presence of a coupling agent such as
 DCC to provide a compound of the formula
 10 XV




A compound of formula XV can then be deprotected with an acid
 e.g., HCl, to provide a compound of the formula
 15 XVI




Compounds of formula XVI are compounds of formula III where V and W
 are $-\text{CH}_2-$, $m=0$, B^2 is aryl, R^1 , R^2 and R^3 are each hydrogen and R is
 20 attached to V and is alkyl having an $-\text{OR}^{11}$ substituent. Compounds of
 formula XVI can be modified as known by those having ordinary skill in
 the art to other compounds of formula III where R is a different alkyl
 group, other than arylsulfonyl or heteroarylsulfonyl substituted alkyl.

- Compounds of formula III where V and W are $-\text{CH}_2-$, $m=0$, B^2 is aryl, R^1 , R^2 and R^3 are each hydrogen, R is attached to V and is an arylsulfonyl or heteroarylsulfonyl substituted alkyl can be prepared by deprotecting compounds of formula XII (with e.g. NCl) to form compounds of formula XVII




- Compounds of formula XVII can be coupled to a compound of formula XIVa in the presence of a coupling agent such as DCC to form compound of formula XVIII




- Treatment of a compound of formula XVIII with an aryldisulfide such as diphenyl disulfide in the presence of a trialkyl phosphine such as tri n-butylphosphine followed by oxidation to the sulfonyl using an oxidizing agent such as the magnesium salt of monoperoxyphthalic acid provides compounds of formula

XIX



A compound of formula XIX can be deprotected (with e.g. HCl) to
 5 provide a compound of formula
 XX



Compounds of formula XX are compounds of formula III where V and W
 are $-\text{CH}_2-$, $m=0$, B^2 is aryl, R^1 , R^2 and R^3 are each hydrogen and R is
 10 attached to V and is an arylsulfonyl substituted alkyl. Compounds of
 formula XX can be modified as known by those having ordinary skill in the
 art to other compounds of formula III where R is a different arylsulfonyl
 or heteroarylsulfonyl substituted alkyl.

Other compounds of formula III can be prepared by modification of
 15 the procedures described herein.

Compounds of formula VII, VIII, XIIa and XIVa can be prepared
 from commercially available materials or are commercially available.

Protecting groups as used herein may be used in the above processes
 with amino acids having reactive functionalities, such as hydroxyl,
 20 carboxyl, amino, mercapto, guanidino, imidazolyl, indolyl and the like.
 The particular protecting groups used for any amino acid residues depend
 upon the sidechains to be protected and are generally known in the art.
 Exemplary sidechain protecting groups include acetyl, benzoyl, benzyl, t-
 butyl and the like for hydroxyl; cyclohexyl, benzyl, methyl, ethyl, t-butyl

and the like for carboxyl; benzyl, 4-methylbenzyl, 4-methoxybenzyl, acetyl, acetamidomethyl, triphenylmethyl (trityl) and the like for mercapto; t-butoxycarbonyl (Boc), benzyloxycarbonyl (Cbz), N-[(9H-Fluoren-9-ylmethoxy)carbonyl] (Fmoc), phthaloyl (Pht), p-toluenesulfonyl (Tos),
5 trifluoroacetyl, 2-(trimethylsilyl)ethoxycarbonyl (Teoc) and the like for amino; 2,4-dinitrophenyl, benzyloxymethyl, Tos, Boc, trityl and the like for imidazolyl; formyl, Cbz, Teoc, 2,2,2-trichloroethyl carbamate (TROC) and the like for indolyl; and tosyl, nitro, bis(1-adamantyloxycarbonyl) and the like for guanidino.

10 Side-chain protecting groups may be removed, if desired, by, for example, treatment with one or more deprotecting agents in an inert solvent or solvent mixture. For examples of protecting groups and suitable deprotecting agents, see M. Bodansky and A. Bodansky, "The Practice of Peptide Synthesis", Springer-Verlag, Inc. (1984); and T. W. Greene and
15 P.G.M. Wuts, "Protective Groups in Organic Synthesis", Second Edition, John Wiley & Sons, New York, 1991.

For additional examples of protecting groups (as well as means of formation and eventual deprotection), see T.W. Greene and P.G.M. Wuts, "Protective Groups in Organic Synthesis", Second Edition, John Wiley &
20 Sons, New York, 1991.


A variety of coupling agents may be used for the coupling of compounds of formula VII with VIII to form compounds of formula IX, including 1-(3-dimethyl-aminopropyl)-3-ethyl-carbodiimide hydrochloride (EDC) with 1-hydroxybenzotriazole (HOBT), dicyclohexyl-carbodiimide
25 (DCC) with HOBT, benzotriazol-1-yloxy-tris (dimethylamino)phosphonium hexafluoro-phosphate (BOP) with or without HOBT, carbonyldiimidazole (CDI), bis(2-oxo-3-oxazolidinyl)phosphinic chloride (BOP chloride), isopropylchloroformate (IPCF), bromo-tris-pyrrolidino-phosphonium hexafluorophosphate (ByBroP) and the like.

30 The following examples and preparations describe the manner and process of making and using the preferred embodiments of the invention and are illustrative rather than limiting. It should be understood that there may be other embodiments which fall within the spirit and scope of the invention as defined by the claims appended hereto.

Example 1

[2S-[2a(R*),4a]]-1-[(4-Mercapto-2-pyrrolidiny)methyl]-2-(2-methoxyethyl)-4-(1-naphthalenylcarbonyl)piperazine,

5 **trifluoroacetate**



A. 1-Benzyl-3(S)-cyclohexyloxycarbonylmethylpiperazine-2,5-dione

10

Boc-L-aspartic acid b-cyclohexyl ester (33 g, 105 mmol) was added to a 0.143 M solution of dicyclohexylcarbodiimide (21.6 g, 105 mmol) in methylene chloride (730 mL) at 0°C. The resulting slurry was stirred for 5 minutes, and ethyl N-benzylglycinate (19.6 mL, 105 mmol) was added.

15

The mixture was stirred for 2 hours at 0°C and at room temperature for 16 hours. The precipitate was removed by filtration, and anhydrous HCl was bubbled through the methylene chloride solution for 6 hours. The solvent was removed *in vacuo*, and the residue partitioned between ethyl acetate (3x300 mL) and 10% sodium bicarbonate (200 mL). The combined organic phases were washed with brine, dried over magnesium sulfate, filtered and evaporated to give Compound A as a white powder (36 g, 100%), mp: 140-143°C.

20

MS: (M+H)⁺ 345.

B. 4-Phenylmethyl-1-[(1,1-dimethylethoxy)carbonyl]-2(S)-(2-hydroxyethyl)piperazine

25

Lithium aluminum hydride (15.1 g, 377 mmol) was added portionwise to a mechanically stirred solution of Compound A (36.0 g, 105 mmol) in THF (780 mL). The reaction mixture was stirred at reflux

for 18 hours, cooled to 0°C, and quenched by the sequential slow addition of water (20 mL), 10% NaOH (20 mL), water (20 mL), and diethylether (100 mL). The mixture was stirred at 0°C for 30 minutes, at room temperature for 30 minutes and filtered. The solvent was removed *in vacuo*, the crude product taken up in methylene chloride (200 mL) and the solution dried over magnesium sulfate. The drying agent was removed by filtration, and the filtrate was treated with di-tert-butyl dicarbonate (25 g, 115 mmol). After 72 hours at room temperature, saturated sodium bicarbonate (300 mL) was added. The layers were separated, the aqueous phase reextracted with methylene chloride (300 mL), and the combined organic extracts dried over magnesium sulfate. Filtration and evaporation gave the crude product which was purified by flash chromatography on silica gel, eluting with 25% ethyl acetate in hexane. Compound B was obtained as a clear viscous oil (20 g, 60%).

MS: (M+H)⁺ 321.

C. 4-Phenylmethyl-1-[(1,1-dimethylethoxy)carbonyl]-2(S)-(2-methoxyethyl)piperazine

Compound B (6.18 g, 19.2 mmol) was dissolved in dry, degassed dimethylformamide and the solution was cooled to 0°C under N₂. Sodium hydride (1.0 g, 24.9 mmol) was added followed by methyl iodide (3.8 g, 1.7 mL). After stirring for 3 hours at 0°C, saturated ammonium chloride was added. The mixture was concentrated under vacuum and the residue was partitioned between ethyl acetate and saturated sodium bicarbonate. The ethyl acetate layer was washed with water, brine and dried over MgSO₄. The crude product was chromatographed (silica gel, 40% ethyl acetate, 60% hexane) to yield Compound C as a clear oil (5.3 g, 82%).

MS: (M+H)⁺ 335.

D. 1-[(1,1-Dimethylethoxy)carbonyl]-2(S)-(2-methoxyethyl)-piperazine

Compound C (4.0 g, 12 mmol) was dissolved in methanol (49 mL) in a Parr bottle and the vessel was purged with Ar. To this was added
5 10% palladium on carbon and the reaction hydrogenated under 60 psi hydrogen for 16 hours. The catalyst was removed by filtration through Celite and the filtrate was concentrated under vacuum to yield Compound D as a clear oil (2.9 g, 99%).

MS: (M+H)⁺ 245.

10

E. 1-[(1,1-Dimethylethoxy)carbonyl]-2(S)-(2-methoxyethyl)- 4-(1-naphthalenylcarbonyl)piperazine

A solution of 1-naphthoic acid (2.35 g, 13.1 mmol), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide•HCl (EDC, 2.74 g, 14.3 mmol), and
15 1-hydroxybenzotriazole (HOBT, 1.93 g, 14.3 mmol) were stirred in DMF (10 mL) at room temperature for 20 minutes. A solution of Compound D (3.0 g, 11.9 mmol) and N,N-diisopropylethylamine (DIEA, 3.5 g, 4.8 mL, 27.4 mmol) in DMF (10 mL) was added dropwise and the mixture was stirred for 16 hours. The mixture was poured into water (200 mL), and
20 extracted with ethyl acetate (3x100 mL). The combined ethyl acetate layers were washed with water (2x100 mL), brine (100 mL), dried over MgSO₄ and concentrated. Chromatography on silica gel (40% ethyl acetate, 60% hexane) yielded Compound E as a slightly brown oil (4.8 g, 99%).

25 MS: (M+H)⁺ 399.

F. 2(S)-(2-Methoxyethyl)-4-(1-naphthalenylcarbonyl)- piperazine, hydrochloride

Compound E (240 mg, 0.60 mmol) was stirred in dimethyl sulfide
30 (0.3 mL) and 4N HCl in dioxane (10 mL) for 30 minutes. The mixture was concentrated, redissolved in methylene chloride (50 mL) and concentrated. This procedure was repeated five times to yield Compound F as a clear glass.

**G. (2S,4S)-[(1-[(1,1-dimethylethoxy)carbonyl]-4-
[(triphenylmethyl)thio]-2-pyrrolidine carboxylic acid methyl ester**

To a solution of (2S,4R)-[(1-[(1,1-dimethylethoxy)-carbonyl]-4-
5 hydroxy-2-pyrrolidine carboxylic acid methyl ester (commercially
available, 1.8 g, 7.3 mmol) and 2,6-di-t-butyl-4-methylpyridine (1.5 g,
7.3 mmol) in methylene chloride (27 mL) at 0°C under nitrogen, was
added triflic anhydride (1.4 mL, 8.1 mmol) in methylene chloride (7 mL)
over 1 hour. The solution was stirred for an additional 15 minutes at 0°C
10 and added directly to a stirred solution of triphenylmethylmercaptan (4.9
g, 18 mmol) and 1M lithium hexamethyldisilazide in THF (18 mL, 18
mmol) in dry THF (30 mL) at 0°C. The mixture was warmed to room
temperature and stirred for 16 hours, concentrated and the residue was
chromatographed (260-400 mesh silica gel, 5.1x15 cm, 1:9-1:3/ethyl
15 acetate:hexane). Fractions containing the desired product were
concentrated to yield 1.5 g of Compound G as a glass. MS (M+H)⁺ 504.

**H. (2S,4S)-[(1-[(1,1-dimethylethoxy)carbonyl]-4-
[(triphenylmethyl)thio]-2-pyrrolidine carboxylic acid**

20 A solution of Compound G (0.50 g, 1.0 mmol) in THF (10 mL) and
methanol (5 mL) was adjusted to pH 12 with 1N LiOH and stirred for 16
hours at room temperature. Ether (50 mL) was added along with water
(30 mL) and the layers were separated. The aqueous layer was washed
with ether (2x40 mL), acidified to pH 3-4 using 1N aqueous KHSO₄ and
25 extracted again with ether (3x50 mL). The combined ether extracts from
the acidic aqueous solution were dried (MgSO₄), pooled and concentrated
to give 0.48 g of Compound H as a glass. MS (M+H)⁺ 489.

**I. (2S,4S)-(N-methoxy-N-methylamino)-[(1-[(1,1-
30 dimethylethoxy)carbonyl]-4-[(triphenylmethyl)thio]-2-
pyrrolidine carboxylic acid amide**

Compound H (0.48 g, 1.0 mmol) was combined with N,O-
dimethylhydroxylamine hydrochloride salt (100 mg, 1.1 mmol), PyBrop

(510 mg, 1.1 mmol) and DIEA (540 μ L, 3.1 mmol) in methylene chloride (30 mL) under argon at room temperature. The mixture was stirred for 2 hours and concentrated under vacuum. The residue was chromatographed (silica gel, 1:3-2:1/ethyl acetate:hexane) and fractions containing the desired compound were concentrated to yield 350 mg of Compound I as a glass. MS (M+H)⁺ 533.

**J. (2S,4S)-[(1-[(1,1-dimethylethoxy)carbonyl]-4-
[(triphenylmethyl)thio]-2-pyrrolidine carboxaldehyde**

Compound I (0.35 g, 0.65 mmol) was dissolved in dry THF (20 mL) and the solution was cooled to 0°C. 1M lithium aluminum hydride in THF (0.66 mL, 0.66 mmol) was added dropwise over 15 minutes. The mixture was stirred for 30 minutes at 0°C under nitrogen. Ether (70 mL) was added, the mixture was quenched with dropwise addition of 1M KHSO₄ (10 mL) at 0°C and stirred for 1 hour at 0°C and the layers were separated. The organic layer was washed with 1M aqueous KHSO₄ (4x50 mL), aqueous saturated sodium bicarbonate (NaHCO₃) (50 mL) and brine (50 mL), dried (MgSO₄), and concentrated to yield Compound J as a white glass (270 mg) which was used immediately in the next step without further purification. R_f 0.25 (ethyl acetate:hexane/1:1, UV, PMA).

**K. [2S-[2a(R*),4a]]-1-[(1-[(1,1-Dimethylethoxy)carbonyl]-4-
triphenylmethylmercapto-2-pyrrolidinyl)methyl]-2-(2-
methoxyethyl)-4-(1-naphthalenylcarbonyl)piperazine**

Compound J (200 mg, 0.60 mmol) and compound F (240 mg, 0.520 mmol) were dissolved in dry methylene chloride (3 mL) and 1,2-dichloroethane (7 mL). Glacial acetic acid (0.5 mL) was added followed by sodium triacetoxyborohydride (160 mg, 0.78 mmol). The mixture was stirred for 16 hours, cooled to 0°C and saturated sodium bicarbonate (2 mL) was added. The mixture was stirred for 20 minutes at 0°C, concentrated and partitioned between ethyl acetate (50 mL) and water (10 mL). The organic layer was separated and the aqueous layer was

extracted with ethyl acetate (2x30 mL). The combined ethyl acetate phases were washed with water (60 mL), brine (50 mL), dried over MgSO_4 and concentrated. The crude product was chromatographed (silica gel, 40% ethyl acetate, 60% hexane) to yield Compound K as a
5 clear oil (90 mg, 23%).
MS $(\text{M}+\text{H})^+$ 756.

L. [2S-[2a(R*),4a]]-1-[(4-Mercapto-2-pyrrolidiny)methyl]-2- (2-methoxyethyl)-4-(1-naphthalenylcarbonyl)piperazine, trifluoroacetate

10 A solution of Compound K (90 mg, 0.12 mmol) in methylene chloride (7 mL), triethylsilane (0.030 mL, 22 mg, 0.19 mmol), and trifluoroacetic acid (7 mL) was stirred for 1 hour at room temperature. The mixture was concentrated, triturated with hexanes and the remaining crude solid was purified by preparative HPLC (YMC S-10
15 ODS column, 30x500 mm; solvent A, 0.1% TFA in 90% water, 10% methanol; solvent B, 0.1% TFA in 10% water, 90% methanol: 20-50% B in 40 minutes, flow rate 20 mL/minute; UV monitored at 220 nm). Fractions containing the desired product were combined, concentrated and lyophilized to provide the compound of Example 1 as a white solid (25
20 mg, 33%), mp 85-86°C.

$[\alpha]_{\text{D}}^{25} = +22^\circ$ ($c=0.01$, MeOH).

MS: $(\text{M}+\text{H})^+$ 414.

Analysis calculated for $\text{C}_{23}\text{H}_{31}\text{N}_3\text{O}_2\text{S} \cdot 2.1 \text{ CF}_3\text{CO}_2\text{H}$.


Calculated: C, 49.54; H, 5.04; N, 6.33.

25 Found: C, 49.09; H, 5.17; N, 6.41.

Example 2

[1a(S),3a]-1-[(3-Mercaptocyclopentyl)carbonyl]-2,2-methoxyethyl-4-(1-naphthalenylcarbonyl)piperazine

5



A. [1a(S),3a]-1-[(3-(Triphenylmethylthio)cyclopentyl)- carbonyl]-2,2-methoxyethyl-4-(1-naphthalenylcarbonyl)- piperazine

10 DIEA (0.23 mL, 1.3 mmol) was added to a solution of (\pm)-cis-3-[(triphenylmethyl)thio]cyclopentanecarboxylic acid [*Can. J. Chem.* **64**, 2184, (1986)] (0.14 g, 0.36 mmol), Compound F of Example 1 (2(S)-(2-methoxyethyl)-4-(1-naphthalenylcarbonyl)- piperazine, hydrochloride; 15 0.13 g, 0.33 mmol), PyBrop (0.23 g, 0.33 mmol) and DMAP (0.61 g, 0.50 mmol) in CH₂Cl₂. The mixture was stirred 16 hours, quenched with 1N HCl (30 mL) and extracted with ethyl acetate (3x60 mL). The combined organic extracts were washed with 10% NaHCO₃ (1x50 mL) and the organic layer was dried (MgSO₄), filtered and concentrated under 20 vacuum. The residue was purified by flash chromatography (eluting with 2/1 hexane/acetone) to afford Compound A (0.19 g, 87%) as a glassy solid, mp: 86-91°C. MS: (M+H)⁺ 669.

B. [1a(S),3a]-1-[(3-Mercaptocyclopentyl)carbonyl]-2,2- methoxyethyl)- 25 4-(1-naphthalenylcarbonyl)piperazine

All solvents used were degassed with argon. Triethylsilane (0.39 mL, 2.6 mmol) was added to a solution of Compound A (0.17 g, 0.26 mmol) in dichloromethane (2 mL). TFA (2 mL) was added and the

mixture was stirred for 3 hours and concentrated under vacuum. The residue was triturated with hexane (3x10 mL) and the remaining residue was purified by preparative HPLC (YMC S10 ODS 30x500 mm, 30-90% aqueous methanol with 0.1% TFA, 60 minute gradient,

- 5 20 mL/minute) and the appropriate fractions were concentrated under vacuum. The residue was dissolved in water (5 mL), millipore filtered and lyophilized to afford the compound of Example 2 (0.04 g, 36%).

MS: (M+H)⁺ 427.

IR (KBr) 2928, 2525, 1674, 1636, 1470, 1437, 1200, 1128 cm⁻¹.

- 10 Analysis calculated for C₂₄H₃₀N₂O₃S•0.2 CF₃CO₂H•0.69 H₂O.


Calculated: C, 63.45; H, 6.89; N, 6.06.

Found: C, 63.45; H, 6.61; N, 6.01.

15

Example 3

[1a(S),3a]-1-[(3-Mercaptocyclopentyl)methyl]-2-(2-methoxyethyl)-4-(1-naphthalenylcarbonyl)piperazine, monohydrochloride



20

A. N,O-Dimethyl-cis-(3-(triphenylmethylthio)cyclopentyl)-hydroxamic acid

- DIEA (1.2 mL, 7.1 mmol) was added to a solution of (±)-cis-3-[(triphenylmethyl)thio]cyclopentanecarboxylic acid (0.93 g, 2.4 mmol),
25 BOP (1.0 g, 2.4 mmol) and N,O-dimethylhydroxylamine HCl (0.24 g, 2.4 mmol) in 3:1 CH₃CN/DMF. The mixture was stirred 16 hours, quenched with 1:1 10% LiCl/1N HCl (100 mL) and extracted with ethyl acetate (3x100 mL). The combined organic extracts were washed with

10% NaHCO₃ (1x50 mL), 10% LiCl (2x100 mL) and the organic layer was dried (MgSO₄), filtered and concentrated under vacuum. The residue was purified by flash chromatography (eluting with 2/1 hexane/acetone) to afford Compound A (0.9 g, 87%) as a glassy solid. MS: (M+H)⁺ 432.

5

B. Cis-3-(triphenylmethylthio)cyclopentanecarboxaldehyde

Lithium aluminum hydride (1M, 2.4 mL, 2.4 mmol) was added to solution of Compound A (0.86 g, 2.0 mmol) in diethyl ether (30 mL) at -15°C. The mixture was stirred at -15°C for 30 minutes and warmed to 0°C and stirred for an additional 1 hour. The mixture was quenched with 10% KHSO₄ (10 mL) and the resulting mixture was stirred 1 hour at room temperature. The mixture was extracted with diethyl ether (3x50 mL). The combined organic extracts were washed with 10% KHSO₄ (1x100 mL), 10% NaHCO₃ (1x100 mL), dried (MgSO₄), filtered and concentrated under vacuum to afford Compound B (0.62 g, 84%) which was used without further purification. MS: (M-H)⁻ 371.

10

15

C. [1a(S),3a]-1-[(3-(Triphenylmethylmercapto)-cyclopentyl)methyl]-2-(2-methoxyethyl)-4-(1-naphthalenylcarbonyl)piperazine

Sodium triacetoxyborohydride (0.35 g, 1.7 mmol) was added to a mixture of Compound B (0.40 g, 1.1 mmol) and Compound F of Example 1 (2(S)-(2-methoxyethyl)-4-(1-naphthalenylcarbonyl)piperazine, hydrochloride) in CH₂Cl₂ (10 mL). The mixture was stirred for 2 hours and quenched with 10% NaHCO₃ (50 mL). The biphasic solution was extracted with CH₂Cl₂ (3x50 mL) and the combined organic extracts were dried (MgSO₄), filtered and concentrated under vacuum. The residue was purified by flash chromatography (eluting with 1/1 hexane/ethyl acetate) to afford Compound C (0.5 g, 80%) as a glassy solid. MS: (M+H)⁺ 655.

20

25

D. [1a(S),3a]-1-[(3-Mercaptocyclopentyl)methyl]-2-(2-methoxyethyl)-4-(1-naphthalenylcarbonyl)piperazine, monohydrochloride

The title compound was prepared from Compound C as described for Compound B of Example 2. Following prep HPLC, the residue was dissolved in methanol (5 mL) and 1N HCl (1 mL) was added. The mixture was stirred 5 minutes and concentrated under vacuum. This latter procedure was repeated two times and the residue was dissolved in water (5 mL), millipore filtered and lyophilized to afford Example 3 as a white solid, mp: 88-98°C.

MS: (M+H)⁺ 413.

Analysis calculated for C₂₄H₃₂N₂O₂S•1.0 HCl•0.88 H₂O.

Calculated: C, 62.00; H, 7.54; N, 6.03.


Found: C, 62.00; H, 7.62; N, 5.96.

15

Example 4

[2S-[2a(R*),4a]]-1-[(4-Mercapto-2-pyrrolidinyl)methyl]-4-(1-naphthalenylcarbonyl)-2-[2-(3-pyridinylmethoxy)ethyl]piperazine, trifluoroacetate

20



25 A. 1-[(1,1-dimethylethoxy)carbonyl]-2(S)-(2-hydroxyethyl)-piperazine

10% Pd/C (0.38 g) was added to a solution of Compound B of Example 1 (4-phenylmethyl-1-[(1,1-dimethylethoxy)carbonyl]-2(S)-(2-

hydroxyethyl)piperazine; 2.1 g, 6.56 mmol) in MeOH (30 mL) at room temperature. The mixture was agitated under hydrogen atmosphere at 53 psi for 20 hours and filtered through a celite pad. The pad was washed 3 times with methanol. The filtrate was concentrated to afford

5 Compound A as a viscous oil (1.58 g, 100%). MS: (M+H)⁺ 231.

B. 1-[(1,1-Dimethylethoxy)carbonyl]-2(S)-(2-hydroxyethyl)-4-(1-naphthalenylcarbonyl)piperazine

Compound B was prepared from Compound A as described for

10 Compound E of Example 1. MS: (M+H)⁺ 385.

C. 1-[(1,1-Dimethylethoxy)carbonyl]-2(S)-[2-((3-pyridinylmethoxy)ethyl)]-4-(1-naphthalenylcarbonyl)-piperazine

15 NaH (0.79 g, 20.8 mmol) was added to a solution of Compound B (0.8 g, 2.0 mmol) in DMF (16 mL) at room temperature under argon. After stirring for 0.5 hour, picolyl chloride (1.64 g, 10 mmol) was added. The mixture was heated at 93°C for 5 hours and the DMF was evaporated. The residue was diluted with saturated NH₄Cl solution (30 mL) and EtOAc (50 mL). The aqueous layer was reextracted with EtOAc (2x50 mL). The combined organic layers were washed with NH₄Cl, NaHCO₃, H₂O, NaCl (3x30 mL), dried over MgSO₄, filtered and concentrated. Flash chromatography (silica, EtOAc) afforded

20 Compound C (0.70 g, 74%). MS: (M+H)⁺ 476.

25

D. [2S-[2a(R*),4a]]-1-[(1-[(1,1-Dimethylethoxy)carbonyl]-4-triphenylmethylmercapto-2-pyrrolidinyl)methyl]-2-((3-pyridinylmethoxy)ethyl)]-4-(1-naphthalenylcarbonyl)-piperazine

30 TFA (5 mL) was added to a solution of Compound C (0.35 g, 0.74 mmol) in CH₂Cl₂ (5 mL). The mixture was stirred 30 minutes and concentrated. The residue was extracted with CHCl₃ (2x15) and the combined extractes were concentrated to afford Compound D.

E. [2S-[2a(R*),4a]]-1-[(1-[(1,1-Dimethylethoxy)carbonyl]-4-triphenylmethylmercapto-2-pyrrolidinyl)methyl]-2-[(3-pyridinylmethoxy)ethyl]-4-(1-naphthalenylcarbonyl)-
5 piperazine

Triethylamine was added to a solution of Compound D in CH₂Cl₂ (2 mL) and the pH was adjusted to 7. A solution of the title J compound of Example 1 ((2S,4S)-[(1-[(1,1-dimethylethoxy)carbonyl]-4-triphenylmethylthio-2-pyrrolidine carboxaldehyde; 0.34 g, 0.72 mmol) in
10 CH₂Cl₂ (2 mL) was added to the above solution, followed by 3A sieves (5 g). The mixture was stirred for 15 minutes under argon. Sodium triacetoxyborohydride (0.63 g, 2.97 mmol) was added. The mixture was stirred for 16 hours and diluted with EtOAc (30 mL) and NaHCO₃ (10 mL). The layers were separated and the aqueous layer was reextracted
15 with EtOAc (2x20 mL). The combined organic layers were combined and washed with NaHCO₃ (20 mL), NaCl (2x20 mL), dried over MgSO₄, filtered and concentrated. Flash chromatography (silica, step gradient of hexane and EtOAc followed by 9/1 CHCl₃/MeOH) afforded Compound E (0.15 g, 26%) as a foamy solid. MS: (M+H)⁺ 833.5.

20


F. [2S-[2a(R*),4a]]-1-[(4-mercapto-2-pyrrolidinyl)methyl]-2-[(3-pyridinylmethoxy)ethyl]-4-(1-naphthalenylcarbonyl)- piperazine

Compound F was prepared from Compound E as described for Compound L of Example 1.

25 MS: (M+H)⁺ 491.

Example 5

[2S-[2a(R*),4B]]-1-[(4-Mercapto-2-pyrrolidiny)methyl]-2-(2-methoxyethyl)-4-(1-naphthalenylcarbonyl)piperazine, dihydrochloride



5

A. (2S,4R)-[(1-[(1,1-dimethylethoxy)carbonyl]-4-[(triphenylmethyl)thio]-2-pyrrolidine carboxylic acid methyl ester

10 Compound A was prepared in a similar manner to Compound G in Example 1 starting with (2S,4R)-[(1-[(1,1-dimethylethoxy)carbonyl]-4-hydroxy-2-pyrrolidine carboxylic acid methyl ester (commercially available). Yield 1.5 g of Compound A as a glass. MS (M+H)⁺ 504.

15 **B. (2S,4R)-[(1-[(1,1-dimethylethoxy)carbonyl]-4-[(triphenylmethyl)thio]-2-pyrrolidine carboxylic acid**

Compound B was prepared in a similar manner to Compound H in Example 1 starting with Compound A. Yield 0.48 g of Compound B as a glass. MS (M+H)⁺ 489.

20

C. (2S,4R)-(N-methoxy-N-methylamino)-[(1-[(1,1-dimethylethoxy)carbonyl]-4-[(triphenylmethyl)thio]-2-pyrrolidine carboxylic acid amide

25 Compound C was prepared in a similar manner to Compound I in Example 1 starting with Compound B. Yield: 350 mg of Compound C as a glass. MS (M+H)⁺ 533.

**D. (2S,4R)-[(1-[(1,1-dimethylethoxy)carbonyl]-4-
[(triphenylmethyl)thio]-2-pyrrolidine carboxaldehyde**

Compound D was prepared in a similar manner to Compound J in Example 1 starting with Compound C. Yield: 270 mg of Compound D
5 as a glassy solid which was used immediately in the next step without further purification. R_f 0.25 (ethyl acetate:hexane/1:1, UV, PMA).

E. [2S-[2a(R*),4b]]-1-[(4-Mercapto-2-pyrrolidinyl)methyl]- 2-(2-methoxyethyl)-4-(1-naphthalenylcarbonyl)- piperazine, dihydrochloride
10

The title compound was prepared from Compound D by the procedure of Compound K of Example 1 followed by the procedure of Compound L of Example 1 except that the HPLC purified compound was dissolved in 1N HCl followed by removal of solvent using lyophilization
15 and this procedure repeated to provide 40 mg (38%) of the compound of Example 5 as a white powder. mp 152-153°C.

MS (M+H)⁺ 414.


Analysis calculated for C₂₃H₃₁N₃O₂S·2.4HCl, 1.1H₂O.

Calculated: C 53.03, H 6.89, N 8.07,

20 Found: C 53.09, H 6.86, N 7.73.

Example 6

**[2R-[2a(S*),4b]]-1-[(4-Mercapto-2-pyrrolidinyl)methyl]-2-(2-methoxyethyl)-
25 4-(1-naphthalenylcarbonyl)piperazine, dihydrochloride**



A. (2R,4R)-[(1-[(1,1-dimethylethoxy)carbonyl]-4-hydroxy-2-pyrrolidine carboxylic acid

A solution of di-t-butyl-di-carbonate (40 g, 0.18 mol) in p-dioxane (200 mL) was added dropwise to a solution of (2R,4R)-4-hydroxy-2-pyrrolidine carboxylic acid (20 g, 0.15 mol) in 10% NaHCO₃ (200 mL) in water over 1 hour and the solution was stirred at 0°C for 20 minute. The solution was allowed to warm to room temperature and stirred for additional 40 hours, concentrated and the resulting aqueous layer was extracted with ethyl acetate (3x300 mL). The organic layers were combined, washed with brine, dried (MgSO₄) and concentrated to yield Compound A as a white solid (20 g, 57%). MS (M+H)⁺ 232.

B. (2R,4R)-[(1-[(1,1-dimethylethoxy)carbonyl]-4-hydroxy-2-pyrrolidine carboxylic acid, methyl ester

To a solution of Compound A (13 g, 56.22 mmol) in methanol (130 mL) was slowly added 2M trimethylsilyl-diazomethane in hexane (37 mL, 73 mmol). The mixture was stirred for 1 hour and concentrated under vacuum to afford 14 g of Compound B. MS (M+H)⁺ 246.

C. [2R-[2a(S*),4b]]-1-[(4-Mercapto-2-pyrrolidinyl)methyl]-2-(2-methoxyethyl)-4-(1-naphthalenylcarbonyl)piperazine, dihydrochloride

The title compound was prepared from Compound B by following the procedures outlined for Compounds G through L of Example 1. The HPLC purified compound was dissolved in 1N HCl followed by removal of solvent using lyophilization. This procedure was repeated to provide the compound of Example 6 as a white powder. mp 144-145°C, MS (M+H)⁺ 414.


Analysis calculated for C₂₃H₃₁N₃O₂S·2.0HCl, 1.4H₂O.

Calculated: C 53.99, H 7.05, N 8.21, Cl 13.86,

Found: C 54.14, H 7.00, N 8.09, 14.14.

Example 7

[2S-[2a(R*),4a]]-1-[(4-Mercapto-2-pyrrolidinyl)methyl]-4-(1-naphthalenylcarbonyl)piperazine, dihydrochloride



5

The title compound was prepared from piperazine using the following procedure: Compound A of Example 6, with chromatography on silica using 1:9-1:3/ethyl acetate:hexane; coupling with 1-naphthoic acid as described for Compound E of Example 1 and Compound F of Example 1; with (2S,4S)-[(1-[(1,1-dimethylethoxy)carbonyl]-4-triphenylmethylmercapto-2-pyrrolidine carboxaldehyde as described in Compound K of Example 1; Compound L of Example 1, except the HPLC purified compound was dissolved in 1N HCl followed by removal of solvent using lyophilization and this procedure was repeated to provide 180 mg (70 %) of the compound of Example 7 as a white powder, mp 189-190°C.

15

MS (M+H)⁺ 356.

Analysis calculated for C₂₀H₂₅N₃OS·2.0HCl, 1.1H₂O:


20 Calculated: C 53.59, H 6.56, N 9.37,

Found: C 53.67, H 6.26, N 9.27,

[α]_D +20.7 (c = 0.6, methanol).

Example 8

[2S-[2a(R*),4a]]-1-[(4-Mercapto-2-pyrrolidinyl)methyl]-4-(1-naphthalenylcarbonyl)-2-[2-(phenylsulfonyl)ethyl]piperazine, trifluoroacetate



A. 1-[(1,1-Dimethylethoxy)carbonyl]-4-(1-naphthalenylcarbonyl)-2(S)-(2-hydroxyethyl)piperazine

A mixture of Compound B of Example 1 (4-phenylmethyl-1[(1,1-dimethylethoxy)carbonyl]-2(S)-(2-hydroxyethyl)piperazine; 1.05 g, 3.28 mmol), Palladium hydroxide (500 mg, 20% on carbon), and methanol (80 mL) was hydrogenated at 60 psi for 18 hours, filtered and concentrated *in vacuo*. The residue was dissolved in DMF. Naphthoic acid (775 mg, 4.5 mmol), EDC (865 mg, 4.5 mmol), 1-hydroxybenzotriazole (607 mg, 4.5 mmol), and triethylamine (1.4 mL, 4.5 mmol) were added, and the mixture was stirred for 18 hours. The mixture was poured into saturated aqueous sodium bicarbonate solution, extracted with ethyl acetate (2x), washed with water and brine, dried, and concentrated. The residue was purified by flash chromatography on silica gel (300 g), eluting with 1:1 hexane: ethyl acetate, to give Compound A as an oil (700 mg, 58%).

B. 1-[(1,1-Dimethylethoxy)carbonyl]-4-(1-naphthalenylcarbonyl)-2(S)-(2-phenylthioethyl)piperazine

To a mixture of Compound A (600 mg, 1.63 mmol) and tetrahydrofuran (5 mL) were added diphenyldisulfide (720 mg, 3.3 mmol)

and tri-n-butyl phosphine (870 μ l, 3.3 mmol). The resulting mixture was stirred for 18 hours and poured into saturated aqueous sodium bicarbonate solution. The solution was extracted with ethyl acetate (2x), washed with 1N sodium hydroxide solution (2x), dried, and concentrated. The residue was combined with similar material from a 0.16 mmol scale reaction and purified by flash chromatography on silica gel (200 g), eluting with 2:1 hexane: ethyl acetate, to Compound B as a clear oil (800 mg, 94%).

10 **C. 1-[(1,1-Dimethylethoxy)carbonyl]-4-(1-naphthalenylcarbonyl)-2(S)-(2-phenylsulfonylethyl)piperazine**

Monoperoxyphthalic acid magnesium salt (2.5 g, 4 mmol) was added to a mixture of Compound B (600 mg, 1.26 mmol) and methanol (5 mL) at 0°C. The mixture was stirred for 2 hours as it slowly warmed to room temperature. Excess sodium thiosulfate solution was added and the mixture was stirred for 10 minutes and poured into saturated aqueous sodium bicarbonate solution, extracted with ethyl acetate (2x), washed with water and brine, dried, and concentrated to give Compound C (510 mg, 80%).

20

D. 4-(1-Naphthalenylcarbonyl)-2(S)-(2-phenylsulfonylethyl)-piperazine

A mixture of Compound C (560 mg, 1.1 mmol), TFA (5 mL), and methylene chloride (5 mL) was stirred at room temperature for 2 hours and concentrated *in vacuo* to give Compound D (210 mg, 37%).

25

E. 1-[(1,1-Dimethylethoxy)carbonyl]-4-(1-naphthalenylcarbonyl)-2(S)-(2-phenylsulfonylethyl)- piperazine

Compound E was prepared from Compound D as outlined for Compound K in Example 1. Yield: 190 mg (52%).

30

F. [2S-[2a(R*),4a]]-1-[(4-Mercapto-2-pyrrolidinyl)methyl]-4- (1-naphthalenylcarbonyl)-2-[2-(phenylsulfonyl)ethyl]-piperazine, trifluoroacetate

5 The title compound was prepared from Compound E as outlined for Compound L in Example 1. Yield: 56 mg of white powder (36%).

MS (M+H)⁺ 524.

Analysis calculated for C₂₈H₃₃N₃O₃S₂·1.2 TFA, 0.6 H₂O:


Calculated: C 54.00; H 5.26; N, 6.16; F, 10.66; S, 9.44,

10 Found: C 54.00; H 5.22, N, 6.05; F, 10.69; S, 9.41.

What is Claimed is:

1. A compound of the formula


I



5

its enantiomers, diastereomers, or pharmaceutically acceptable salts, prodrugs or solvates thereof where

A is



10 optionally substituted, at any available position or positions on the ring, with one or more of the following substituents, halo, alkyl, alkoxy, aryl, hydroxy, alkanoyl, alkanoyloxy, amino, alkylamino, dialkylamino, alkanoylamino, thiol, alkylthio, alkylthiono, alkylsulfonyl, sulfonamido, nitro, cyano, carboxy, carbamyl or alkoxycarbonyl;

15 B is $-\text{CH}_2-$ or $-\text{C}(\text{O})-$;

B^2 is alkyl, aryl or heterocycle;

V and W are each, independently, $-\text{CH}_2-$ or $-\text{C}(\text{O})-$;

X is $-\text{SH}$, $-\text{OH}$ or $-\text{NHR}^6$;

X' is $-\text{NR}^7-$, $-\text{CH}_2-$ or $-\text{CH}(\text{NHR}^8)-$;

20 R, R^1 , R^2 and R^3 are each independently, hydrogen, alkyl, alkenyl, alkynyl, aryl, heterocycle, $-\text{C}(\text{O})\text{NR}^4\text{R}^5$, or $-\text{C}(\text{O})\text{OR}^4$; or any two of R, R^1 , R^2 and R^3 , if present, may also be alkylene attached to the same carbon atom forming a spiro ring;

R^4 and R^5 are each independently selected from hydrogen,

25 hydroxyl, alkyl, cycloalkyl, heterocycle, aryl, aroyl, heteroaroyl, arylsulfonyl or heteroarylsulfonyl; or R^4 and R^5 may together form a 5- to 7-membered saturated ring with the atom to which they are attached;

said ring being optionally substituted with halogen, hydroxy, alkoxy, oxo, nitro, cyano, $-C(O)H$, $-C(O)OH$ and the like;


R^6 , R^7 and R^8 are each independently, hydrogen or alkyl;

m is an integer from 0 to 1; and

5 n is an integer from 1 to 2.

2. A compound of Claim 1 wherein

A is



10 B is $-CH_2-$ or $-C(O)-$;

B^2 is aryl;

V and W are each, independently, $-CH_2-$;


X is $-SH$;

X' is $-NR^7-$ or $-CH_2-$; and

15 R^7 is hydrogen or alkyl.

3. A compound of Claim 1, wherein

A is



20 B is $-CH_2-$;

B^2 is aryl;

V and W are each, independently, $-CH_2-$;


X is $-SH$;

X' is $-NR^7-$ or $-CH_2-$; and

25 R^7 is hydrogen or alkyl.

4. A compound of Claim 1 wherein


A is



- 5 B is -C(O)-;
 B² is aryl;
 V and W are each, independently, -CH₂-;
 X is -SH;
 X' is -NR⁷- or -CH₂-; and
 10 R⁷ is hydrogen or alkyl.

5. A compound of Claim 1 wherein


A is



- 15 B is -CH₂- or -C(O)-;
 B² is aryl;
 V and W are each, independently, -CH₂-;
 X is -SH;
 X' is -NR⁷- or -CH₂-;
 20 R is alkyl;
 R¹, R² and R³ are each hydrogen;
 R⁷ is hydrogen or alkyl;
 m is 0; and
 n is 1.

6. A compound of Claim 1 wherein

A is



5 B is $-\text{CH}_2-$;

B^2 is aryl;

V and W are each, independently, $-\text{CH}_2-$;

X is $-\text{SH}$;

X' is $-\text{NR}^7-$ or $-\text{CH}_2-$;

10 R is alkyl;

R^1 , R^2 and R^3 are each hydrogen;

R^7 is hydrogen or alkyl;


m is 0; and

n is 1.

15

7. A compound of Claim 1 wherein

A is



B is $-\text{C}(\text{O})-$;

20 B^2 is aryl;

V and W are each, independently, $-\text{CH}_2-$;

X is $-\text{SH}$;

X' is $-\text{NR}^7-$ or $-\text{CH}_2-$;

R is alkyl;

25 R^1 , R^2 and R^3 are each hydrogen;


R^7 is hydrogen or alkyl;

m is 0; and

n is 1.

8. A compound of Claim 1 wherein

A is



B is $-\text{CH}_2-$ or $-\text{C}(\text{O})-$;

5 B^2 is aryl;

V and W are each, independently, $-\text{CH}_2-$;

X is $-\text{SH}$;

X' is $-\text{NR}^7-$ or $-\text{CH}_2-$;

R is $-\text{OR}^{11}$ substituted alkyl;

10 R^1 , R^2 and R^3 are each hydrogen;


R^7 is hydrogen or alkyl;

m is 0; and

n is 1.

15 9. A compound of Claim 1 wherein

A is



B is $-\text{CH}_2-$ or $-\text{C}(\text{O})-$;

B^2 is aryl;

20 V and W are each, independently, $-\text{CH}_2-$;

X is $-\text{SH}$;

X' is $-\text{NR}^7-$ or $-\text{CH}_2-$;

R is arylsulfonyl substituted alkyl;

R^1 , R^2 and R^3 are each hydrogen;


25 R^7 is hydrogen or alkyl;

m is 0; and

n is 1.

10. A compound of Claim 1 wherein

A is



5 B is $-\text{CH}_2-$ or $-\text{C}(\text{O})-$;

B^2 is aryl;

V and W are each, independently, $-\text{CH}_2-$;

X is $-\text{SH}$;

X' is $-\text{NR}^7-$ or $-\text{CH}_2-$;

10 R is arylsulfonyl substituted alkyl;

R^1 , R^2 and R^3 are each hydrogen;

R^7 is hydrogen or alkyl;

m is 0; and

n is 1.

15

11. A compound of Claim 1 which is

[2S-[2a(R*),4a]]-1-[(4-mercapto-2-pyrrolidiny)methyl]-2-(2-methoxyethyl)-4-(1-naphthalenylcarbonyl)piperazine, trifluoroacetate;

20 [1a(S),3a]-1-[(3-mercaptocyclopentyl)carbonyl]-2,2-methoxyethyl)-4-(1-naphthalenylcarbonyl)piperazine;

[1a(S),3a]-1-[(3-mercaptocyclopentyl)methyl]-2-(2-methoxyethyl)-4-(1-naphthalenylcarbonyl)piperazine, monohydrochloride;

25 [2S-[2a(R*),4a]]-1-[(4-mercapto-2-pyrrolidiny)methyl]-4-(1-naphthalenylcarbonyl)-2-[2-(3-pyridinylmethoxy)ethyl]-piperazine, trifluoroacetate;

[2S-[2a(R*),4β]]-1-[(4-mercapto-2-pyrrolidiny)methyl]-2-(2-methoxyethyl)-4-(1-naphthalenylcarbonyl)piperazine, dihydrochloride;

30 [2R-[2a(S*),4β]]-1-[(4-mercapto-2-pyrrolidiny)methyl]-2-(2-methoxyethyl)-4-(1-naphthalenylcarbonyl)piperazine, dihydrochloride;

[2S-[2a(R*),4a]]-1-[(4-mercapto-2-pyrrolidinyl)methyl]-4-(1-naphthalenylcarbonyl)piperazine, dihydrochloride;

[2S-[2a(R*),4a]]-1-[(4-mercapto-2-pyrrolidinyl)methyl]-4-(1-naphthalenylcarbonyl)-2-[2-(phenylsulfonyl)ethyl]-piperazine,

5 trifluoroacetate; or

a pharmaceutically acceptable salt thereof.

12. A method of inhibiting farnesyl protein transferase which comprises administering to a mammalian subject an effective farnesyl protein
10 transferase inhibiting amount of a compound of Claim 1.

13. A method of inhibiting prenyl transferases which comprises administering to a mammalian subject an effective prenyl transferase inhibiting amount of a compound of Claim 1.

15

14. A method of inhibiting tumors which comprises administering to a mammalian subject an effective tumor inhibiting amount of a compound of Claim 1.

20 15. A method of treating diseases associated with signal transduction pathways operating through Ras which comprises administering to a mammalian subject an amount of a compound of Claim 1 effective for treating said diseases.

25 16. A method of treating diseases associated with proteins that are post-translationally modified by the enzyme farnesyl protein transferase which comprises administering to a mammalian subject an amount of a compound of Claim 1 effective for treating said diseases.

30 17. A method of treating diseases associated with proteins that are post-translationally modified by the enzymes geranylgeranyl protein transferase which comprises administering to a mammalian subject an amount of a compound of Claim 1 effective for treating said diseases.

18. A method of treating autoimmune diseases which comprises administering to a mammalian subject an amount of a compound of Claim 1 effective for treating said diseases.

5

19. A pharmaceutical composition comprising a pharmaceutical carrier and a therapeutically effective amount of a compound of Claim 1.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US97/20020

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :C07D 241/04, 243/08, 295/192, 403/06; A61K 31/495, 31/55

US CL :514/218, 252, 255; 544/372, 387, 391; 540/492, 575

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/218, 252, 255; 544/372, 387, 391; 540/492, 575

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAS ONLINE STRUCTURE SEARCH

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	Chem. Abstr. Vol. 123, No. 21, 20 November 1995 (Columbus, OH, USA), page 1213, column 1, the abstract no. 286083j, MATSUO, M. 'Preparation of piperazine-derivative tachykinin antagonists', abstract, EP 655442, 31 May 1995. See pyrrolidine derivative indexed in Vol. 123 Formula Index.	1-11, 19
A	Chem. Abstr., Vol. 122, No. 7, 13 February 1995 (Columbus, OH, USA), page 1049, column 1, the abstract no. 80945n, SUNAGAWA, M. 'Novel quaternary ammonium carbapenems: 1 β -methyl-2-(5'-substituted pyrrolidinylthio)carbapenems', abstract, J. Antibiot. 1994, 47(11), 1337-1340 (Eng.).	1-11, 19

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T*	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A*	document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means		
P document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

16 JANUARY 1998

Date of mailing of the international search report

10 FEB 1998

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

EMILY BERNHARDT

Telephone No. (703) 308-1235